

Some Remarks about the Two-Matrix Penner Model and the Kazakov–Migdal Model

Yu. Makeenko¹

The Niels Bohr Institute, 2100 Copenhagen, DK

and

Institute of Theoretical and Experimental Physics, 117259 Moscow, RF

Abstract

I consider the Hermitean two-matrix model with a logarithmic potential which is associated in the one-matrix case with the Penner model. Using loop equations I find an explicit solution of the model at large N (or in the spherical approximation) and demonstrate that it solves the corresponding Riemann–Hilbert problem. I construct the potential of the Kazakov–Migdal model on a D -dimensional lattice, which turns out to be a sum of two logarithms as well, whose large- N solution is given by the same formulas. In the “naive” continuum limit this potential recovers in $D < 4$ dimensions the standard scalar theory with quartic self-interaction. I exploit the solution to calculate explicitly the pair correlator of gauge fields in the Kazakov–Migdal model with the logarithmic potential.

Submitted to *Physics Letters B*

¹E-mail: makeenko@nbivax.nbi.dk / makeenko@desyvax.bitnet / makeenko@vxitep.itep.msk.su

1 Introduction

The Penner model [1] is the Hermitean *one-matrix* model defined by the partition function

$$Z_1 = \int d\phi e^{-N \operatorname{tr} V(\phi)} \quad (1.1)$$

with the logarithmic potential

$$V(\phi) = -\alpha \left(\log(1 + \phi) - \phi \right). \quad (1.2)$$

This model has been extensively studied in the modern context of matrix models after the observation by Distler and Vafa [2] that the partition function (1.1) with the potential (1.2) coincides with that of the compactified one-dimensional string at a self-dual radius. Some generalizations of the Penner model [3] were considered and solved in the large- N limit. Of special interest is the fact [4] that an external field problem for the Penner model is equivalent to the Hermitean one-matrix model (1.1) with a general potential $V(\phi)$ which is analytic at $\phi = 0$.

In the present paper I consider the Hermitean *two-matrix* model

$$Z_2 = \int d\phi_1 d\phi_2 e^{N \operatorname{tr} \left(\phi_1 \phi_2 - \mathcal{V}(\phi_1) - \mathcal{V}(\phi_2) \right)} \quad (1.3)$$

with the Penner potential (1.2). One of the motivations is the recent paper [5] where the two-dimensional minimal conformal field theories have been constructed from the two-matrix model (1.3) with polynomial potentials as was first conjectured by Douglas [6]. From this point of view it is interesting to find out what happens for the *non-polynomial* potential (1.2).

Another motivation is that the two-matrix model (1.3) describes the solution of the Kazakov–Migdal [7] model which is defined by the partition function

$$Z_{KM} = \int \prod_{x,\mu} dU_\mu(x) \prod_x d\phi_x e^{\sum_x N \operatorname{tr} \left(-V(\phi_x) + \sum_{\mu=1}^D \phi_x U_\mu(x) \phi_{x+\mu} U_\mu^\dagger(x) \right)}, \quad (1.4)$$

where the integration over the gauge field $U_\mu(x)$ is over the Haar measure on $SU(N)$ at each link of a D -dimensional lattice with x labeling its sites. The Kazakov–Migdal model is an interesting extension of the matrix models (1.1) and (1.3) to the multi-dimensional case and obviously recovers the two-matrix model if the lattice is just two-points since the gauge field can be absorbed by a unitary transformation, say of ϕ_1 .

While the fact that the models (1.3) and (1.4) are equivalent at large N is known [8], an explicit relation between the potentials $\mathcal{V}$ in (1.3) and V in (1.4) has been studied recently [9]. The idea of the DMS approach is first to solve the two-matrix model (1.3) in the large- N limit for some choice of $\mathcal{V}$ which is practically convenient and then, having the solution, to calculate the associated potential V of the Kazakov–Migdal model. To realize this program, the loop equation for the one-link correlator was derived whose solution

completely determines all the correlators of the model, in particular, the pair correlator of the gauge fields. Besides the Gaussian case when both potentials are quadratic, some explicit relations between $\mathcal{V}$ and V have been studied.

In the present paper I show that for the Penner potential (1.2) (in fact for a little bit more general one given by Eq. (4.1) below) the associated potential V of the Kazakov–Migdal model is given by the sum of two logarithms as well (Eq. (4.19) below) so that the model (1.4) with this potential is explicitly solved at large N for any D . The solution is conveniently described via the one-matrix model whose potential (Eq. (4.9) below) coincides with V of the Kazakov–Migdal model for $D = 0$. I demonstrate that this solution satisfies the master field equation derived by Migdal [10] and recovers by turning the parameters of the potential the solution for the quadratic potential found by Gross [11]. I study some properties of the Kazakov–Migdal model with this logarithmic potential and show that in the “naive” continuum limit it reproduces in $D < 4$ the standard action of the scalar matrix field with quartic self-interaction. I calculate explicitly the correlator of two gauge fields $U_\mu(x)$ and $U_\mu^\dagger(x)$ at the same link (x, μ) which completely determines all other correlators of the model and show that it might undergo phase transitions at the values of parameters where the proper one-matrix model does.

2 A review of the DMS approach

Let us define for the Kazakov–Migdal model (1.4) the loop average

$$E_\lambda \equiv \left\langle \frac{\text{tr}}{N} \left(\frac{1}{\lambda - \phi_x} \right) \right\rangle \quad (2.1)$$

which describes all correlators of powers of ϕ_x at the same site and the one-link correlator

$$G_{\nu\lambda} \equiv \left\langle \frac{\text{tr}}{N} \left(\frac{1}{\nu - \phi_x} U_\mu(x) \frac{1}{\lambda - \phi_{x+\mu}} U_\mu^\dagger(x) \right) \right\rangle. \quad (2.2)$$

Both functions are analytic on the complex plane with some cut (or cuts) at the real axis and obey the following asymptotic expansions

$$E_\nu = \frac{1}{\nu} + \sum_{k=1}^{\infty} \frac{\mathcal{E}_k}{\nu^{k+1}} \quad , \quad \mathcal{E}_k = \left\langle \frac{\text{tr}}{N} \phi_x^k \right\rangle \quad (2.3)$$

and

$$G_{\nu\lambda} = \frac{E_\lambda}{\nu} + \sum_{n=1}^{\infty} \frac{G_n(\lambda)}{\nu^{n+1}} \quad , \quad G_n(\lambda) = \left\langle \frac{\text{tr}}{N} \left(\phi_x^n U_\mu(x) \frac{1}{\lambda - \phi_{x+\mu}} U_\mu^\dagger(x) \right) \right\rangle. \quad (2.4)$$

Notice that $G_{\nu\lambda}$ is symmetric in ν and λ due to invariance of the Haar measure, dU , under transformations $U \rightarrow U^\dagger$.

The correlator $G_{\nu\lambda}$ obeys in the large- N limit the following equation [9]

$$\int_{C_1} \frac{d\omega}{2\pi i} \frac{\mathcal{V}'(\omega)}{\nu - \omega} G_{\omega\lambda} = E_\nu G_{\nu\lambda} + \lambda G_{\nu\lambda} - E_\nu, \quad (2.5)$$

where the contour C_1 encircles counterclockwise the cut (or cuts) of the function $G_{\nu\lambda}$ (the same as the support of ρ) and

$$\mathcal{V}'(\omega) \equiv V'(\omega) - (2D - 1)F(\omega). \quad (2.6)$$

The function

$$F(\omega) = \sum_{n=0}^{\infty} F_n \omega^n \quad (2.7)$$

is the one which appears in the expansion of the pair correlator of the gauge fields

$$\left\langle \frac{1}{N} \text{tr} (t^a U \chi U^\dagger) \right\rangle_U \equiv \frac{\int dU e^{N \text{tr} (\phi U \chi U^\dagger)} \frac{1}{N} \text{tr} (t^a U \chi U^\dagger)}{\int dU e^{N \text{tr} (\phi U \chi U^\dagger)}} \quad (2.8)$$

where the averaging is only w.r.t. U while ϕ and χ play the role of external fields. As was proposed in Refs. [10, 12], the following formula holds at $N = \infty$:

$$\left\langle \frac{\text{tr}}{N} (t^a U \chi U^\dagger) \right\rangle_U = \sum_{n=1}^{\infty} F_n \frac{\text{tr}}{N} (t^a \phi^n), \quad (2.9)$$

where t^a ($a = 1, \dots, N^2 - 1$) stand for the generators of the $SU(N)$ which are normalized by

$$\sum_a (t^a)^{ij} (t^a)^{kl} = N \delta^{il} \delta^{kj} - \delta^{ij} \delta^{kl}. \quad (2.10)$$

The structure of the r.h.s. of Eq. (2.9) can be understood analyzing the power series expansion of the Itzykson–Zuber correlator in ϕ and χ :

$$\left\langle \frac{\text{tr}}{N} (t^a U t^b U^\dagger) \right\rangle_U = \sum_{n,m} C_{nm} \frac{\text{tr}}{N} (t^a \phi^n) \frac{\text{tr}}{N} (t^b \chi^m), \quad (2.11)$$

where the coefficients C_{nm} depend only on the distribution of eigenvalues of ϕ and χ . This formula is due to the invariance of the Haar measure dU , which forbids terms like $\text{tr} \phi \chi$, as well as to the fact that only the traceless parts of ϕ and χ contribute to the integral in Eq. (2.11). Multiplying Eq. (2.11) by $\text{tr} t^b \chi$ and using the completeness condition (2.10), one identifies F_n with

$$F_n = \sum_m C_{n,m} \left(\frac{\text{tr}}{N} \chi^{m+1} - \frac{\text{tr}}{N} \chi \frac{\text{tr}}{N} \chi^m \right), \quad \text{for } n \geq 1. \quad (2.12)$$

Eq. (2.6) arise after the multiplication of Eq. (2.9) by $\text{tr} (t^a / (\nu - \phi))$ which gives

$$\left\langle \frac{\text{tr}}{N} \left(\frac{1}{\nu - \phi} U \chi U^\dagger \right) \right\rangle_U = \frac{\text{tr}}{N} \left(\frac{F(\phi)}{\nu - \phi} \right) + \frac{\text{tr}}{N} \left(\frac{1}{\nu - \phi} \right) \frac{\text{tr}}{N} (\phi - F(\phi)). \quad (2.13)$$

Since F_0 does not enter this formula, one can always choose

$$F_0 = \frac{\text{tr}}{N} \left(\phi - \sum_{n=1}^{\infty} F_n \phi^n \right) \quad (2.14)$$

to provide

$$\frac{\text{tr}}{N} \left(\phi - F(\phi) \right) = 0. \quad (2.15)$$

For the even potential $V(\phi) = V(-\phi)$ Eq. (2.15) holds automatically since the distribution of eigenvalues is symmetric, $\rho(\lambda) = \rho(-\lambda)$, and $F(\lambda) = -F(-\lambda)$.

With F_0 given by (2.14), Eq. (2.13) can be rewritten as

$$G_1(\nu) = \int_{C_1} \frac{d\omega}{2\pi i} \frac{F(\omega)}{\nu - \omega} E_\omega \quad (2.16)$$

where $G_1(\nu)$ is defined by the expansion (2.4). The same quantity appears in the simplest loop equation (see [13] for a review)

$$\int_{C_1} \frac{d\omega}{2\pi i} \frac{V'(\omega)}{\nu - \omega} E_\omega - 2DG_1(\nu) = E_\nu^2 \quad (2.17)$$

which is just a result of the infinitesimal shift

$$\phi_x \rightarrow \phi_x + \frac{\epsilon}{\nu - \phi_x} \quad (2.18)$$

of the measure in (1.4) and can be alternatively obtained taking the $1/\lambda$ term of the expansion of Eq. (2.5) in λ . Inserting (2.16) into Eq. (2.17) and using (2.6), one arrives at the following equation for E_ν

$$\int_{C_1} \frac{d\omega}{2\pi i} \frac{\tilde{V}'(\omega)}{\nu - \omega} E_\omega = E_\nu^2 \quad (2.19)$$

where

$$\tilde{V}'(\lambda) = \mathcal{V}'(\lambda) - F(\lambda). \quad (2.20)$$

One can see now that Eq. (2.15) can be rewritten using Eqs. (2.19) and (2.20) as

$$\int_{C_1} \frac{d\omega}{2\pi i} \mathcal{V}'(\omega) E_\omega = \mathcal{E}_1 \quad (2.21)$$

which coincides with the $1/(\nu\lambda)$ term of the double expansion of Eq. (2.5) in $1/\nu$ and $1/\lambda$. Therefore, the choice (2.14) is always compatible with Eq. (2.5).

While Eq. (2.19) coincides with the large- N loop equation for the Hermitean one-matrix model, the potential $\tilde{V}(\omega)$ is, generally speaking, non-polynomial and has singularities on the complex plane outside of the cut (or cuts) of E_ω . Another interesting property of the above formulas is that at $D = 1/2$, which is associated with the Hermitean two-matrix model, the last two terms on the r.h.s. of Eq. (2.6) disappear and one gets just $\mathcal{V}(\omega) = V(\omega)$. For this reason the Schwinger–Dyson equations for the Hermitean two-matrix model [14, 15] are equivalent to Eq. (2.5).

3 Exact solution for arbitrary potential

Let us consider the two-matrix model with an arbitrary potential

$$\mathcal{V}(\omega) = \sum_{m=1}^{\infty} \frac{g_m}{m} \omega^m. \quad (3.1)$$

The solution of loop equations in this case was obtained in Ref. [14] and in a more explicit form in Ref. [9]. The equation (2.5) for the potential (3.1) reads explicitly

$$(\lambda + E_\nu - \mathcal{V}'(\nu)) G_{\nu\lambda} = E_\nu - R_\lambda(\nu) \quad (3.2)$$

where $R_\lambda(\nu)$ is given by

$$R_\lambda(\nu) \equiv - \int_{C_2} \frac{d\omega}{2\pi i} \frac{\mathcal{V}'(\omega)}{\nu - \omega} G_{\omega\lambda} = \sum_{m=2}^{\infty} g_m \sum_{n=0}^{m-2} G_n(\lambda) \nu^{m-n-2} \quad (3.3)$$

and the contour C_2 encircles both singularities of $G_{\omega\lambda}$ and the pole at $\omega = \nu$. The terms on the r.h.s. result from taking the residue at infinity in the contour integral and the functions $G_n(\lambda)$ are defined by Eq. (2.4).

The formal solution to Eq. (3.2) is

$$G_{\nu\lambda} = \frac{E_\nu - R_\lambda(\nu)}{\lambda + E_\nu - \mathcal{V}'(\nu)} \quad (3.4)$$

with $R_\lambda(\nu)$ given by (3.3). The functions $G_n(\lambda)$ can be expressed via E_λ using the recurrence relation

$$G_{n+1}(\lambda) = \int_{C_1} \frac{d\omega}{2\pi i} \frac{\mathcal{V}'(\omega)}{\lambda - \omega} G_n(\omega) - E_\lambda G_n(\lambda), \quad G_0(\lambda) = E_\lambda \quad (3.5)$$

which can be obtained expanding Eq. (2.5) in $1/\lambda$. For $n = 0$ this equation recovers Eq. (2.16).

To determine E_λ , I use the equation [9]

$$\int_{C_1} \frac{d\omega}{2\pi i} \mathcal{V}'(\omega) G_{\omega\lambda} = \lambda E_\lambda - 1 \quad (3.6)$$

which is nothing but the $1/\nu$ term of the expansion of Eq. (2.5) in $1/\nu$. Using the expansion (2.4), one can rewrite Eq. (3.6) as

$$\sum_{m \geq 1} g_m G_{m-1}(\lambda) = \lambda E_\lambda - 1. \quad (3.7)$$

For $\mathcal{V}(\lambda)$ being a polynomial of the highest power J , Eq. (3.7) contains E_λ up to the power J and the solution is therefore *algebraic*. This fact was first noted by Staudacher [14] for the two-matrix model. An explicit example of a non-polynomial potential which results in a quadratic equation for E_λ is given below.

As is proven in Ref. [9]:

- Equations which appear from the next terms of the $1/\nu$ -expansion of Eq. (2.5) are automatically satisfied as a consequence of Eqs. (3.5) and (3.7).
- $G_{\nu\lambda}$ given by (3.4) is symmetric in ν and λ for any solution of Eq. (3.7). The symmetry requirement can be used directly to determine E_λ alternatively to Eq. (3.7).

It is worth mentioning the relation to the approach by Migdal [10] which is based on the Riemann–Hilbert method. As is noticed in Ref. [9], Eq. (2.5) is equivalent to the equation of Ref. [10]. To show this let us define the continuous and discontinuous in ν parts of $G_{\nu\lambda}$ across the cut (cuts) by

$$\begin{aligned}\text{Disc}_\nu G_{\nu\lambda} &\equiv \frac{G_{(\nu+i0)\lambda} - G_{(\nu-i0)\lambda}}{2}, \\ \text{Cont}_\nu G_{\nu\lambda} &\equiv \frac{G_{(\nu+i0)\lambda} + G_{(\nu-i0)\lambda}}{2}\end{aligned}\quad (3.8)$$

so that for a real λ outside of the cut (cuts) $\text{Disc}_\nu G_{\nu\lambda}$ coincides with the imaginary part and $\text{Cont}_\nu G_{\nu\lambda}$ coincides with the real part. In particular, $\text{Disc}_\nu E_\nu = -\pi\rho(\nu)$. The discontinuous part of Eq. (2.5) then reads

$$\text{Cont}_\nu G_{\nu\lambda} = 1 + \frac{1}{\pi\rho(\nu)}\left(\lambda + \frac{1}{2}\tilde{V}'(\nu) - \mathcal{V}'(\nu)\right)\text{Disc}_\nu G_{\nu\lambda}, \quad \text{for } \nu \in \text{cut} \quad (3.9)$$

which coincides with the equation of Ref. [10].

To obtain the solution to Eq. (3.9) for $G_{\nu\lambda}$ versus E_ν , one notices that for any real ν

$$\frac{1 - G_{(\nu+i0)\lambda}}{1 - G_{(\nu-i0)\lambda}} = \frac{1 - \text{Cont}_\nu G_{\nu\lambda} - i \text{Disc}_\nu G_{\nu\lambda}}{1 - \text{Cont}_\nu G_{\nu\lambda} + i \text{Disc}_\nu G_{\nu\lambda}} = \frac{\lambda + \text{Re } E_\nu - i \text{Im } E_\nu - \mathcal{V}'(\nu)}{\lambda + \text{Re } E_\nu + i \text{Im } E_\nu - \mathcal{V}'(\nu)} \quad (3.10)$$

since $\text{Disc}_\nu G_{\nu\lambda}$ cancels at the cut (cuts) due to Eq. (3.9). The analytic function which solves this problem is given by [10, 11, 16]

$$G_{\nu\lambda} = 1 - \exp\left\{-\int_{C_1} \frac{d\omega}{2\pi i} \frac{1}{\nu - \omega} \log(\lambda + E_\omega - \mathcal{V}'(\omega))\right\}. \quad (3.11)$$

Taking the residue at $\omega = \nu$, one can rewrite it in the form (3.4) with

$$R_\lambda(\nu) = \exp\left\{-\int_{C_2} \frac{d\omega}{2\pi i} \frac{1}{\nu - \omega} \log\left(1 + \frac{E_\omega}{\lambda - \mathcal{V}'(\omega)}\right)\right\}. \quad (3.12)$$

The l.h.s. is obviously analytic in λ , expandable in $1/\lambda$, has no discontinuity in ν and is expandable in powers of ν — *i.e.* has all the properties required for $R_\lambda(\nu)$.

The formula (3.11) should solve the recurrence relation (3.5). I have checked this expanding in $1/\lambda$ to few lower orders. Analogously, the master field equation

$$E_\lambda = \int_{C_1} \frac{d\omega}{2\pi i} \log(\lambda + E_\omega - \mathcal{V}'(\omega)), \quad (3.13)$$

which is obtained as the $1/\nu$ term of Eq. (3.11), should be equivalent to Eq. (3.7). It is a matter of practical convenience which form of equations to solve. As was shown in Ref. [9], Eqs. (2.5) and (3.7) can easily be solved for the quadratic potential $\mathcal{V}(\phi)$. A simple solution of Eqs. (2.5), (3.7) for the Penner potential is obtained in the next section.

4 An explicit solution for Penner potential

Let us choose the following potential of the two matrix model

$$\mathcal{V}(\phi) = -(ab + c) \log(b - \phi) - a\phi = \sum_{m=2}^{\infty} \frac{ab + c}{m b^m} \phi^m + \frac{c}{b} \phi \quad (4.1)$$

where a and b are real. This potential recovers the quadratic one in the limit

$$a, b \rightarrow \infty, \quad \frac{a}{b} \equiv \frac{1}{\Lambda} \sim 1, \quad c \sim 1 \quad (\text{quadratic potential}). \quad (4.2)$$

The loop equation (2.5) involves

$$\mathcal{V}'(\omega) = \frac{a\omega + c}{b - \omega}. \quad (4.3)$$

The function $R_\lambda(\nu)$ which enters (3.4) can then be calculated to give

$$R_\lambda(\nu) \equiv - \int_{C_2} \frac{d\omega}{2\pi i} \frac{\mathcal{V}'(\omega)}{\nu - \omega} G_{\omega\lambda} = \frac{ab + c}{b - \nu} G_{b\lambda}. \quad (4.4)$$

This formula is obtained doing the integration over the contour C_2 which encircles both singularities of $G_{\omega\lambda}$ and the pole at $\omega = \nu$. For $\mathcal{V}'(\omega)$ given by (4.3) the residue at infinity vanishes since $G_{\omega\lambda}$ decreases as $1/\omega$ while the residue at $\omega = b$ results in the expression on the r.h.s.. One can alternatively derive this formula expanding $\mathcal{V}'(\omega)$ in ω and taking the residue at infinity.

The function $G_{b\lambda}$ on the r.h.s. of Eq. (4.4) can be determined from Eq. (3.6) to be

$$G_{b\lambda} = \frac{(a + \lambda)E_\lambda - 1}{ab + c} \quad (4.5)$$

so that the solution to Eq. (2.5) for $G_{\nu\lambda}$ versus E_λ is

$$G_{\nu\lambda} = \frac{E_\nu - \frac{(a+\lambda)E_\lambda - 1}{b-\nu}}{\lambda + E_\nu - \frac{a\nu+c}{b-\nu}}. \quad (4.6)$$

To determine E_λ let us utilize the symmetry of $G_{\nu\lambda}$ in ν and λ . From Eq. (4.6) one gets

$$G_{\nu b} = \frac{(b - \nu)E_\nu - (a + b)E_b + 1}{(b - \nu)(b + E_\nu) - a\nu - c} \stackrel{(4.5)}{=} \frac{(a + \nu)E_\nu - 1}{ab + c} \quad (4.7)$$

which yields the *quadratic* equation for E_ν

$$\begin{aligned} (a + \nu)(b - \nu)E_\nu^2 - \left((a\nu + c)(a + \nu) + (1 + c - b\nu)(b - \nu) \right) E_\nu \\ = -\nu(a + b) + b^2 - c - (ab + c)((a + b)E_b - 1). \end{aligned} \quad (4.8)$$

I have verified that (4.6) is indeed symmetric in ν and λ providing Eq. (4.8) is satisfied. The formula for $C(\nu, \lambda)$, the double discontinuity of $G_{\nu\lambda}$ both in ν and λ across the cut (cuts), which is manifestly symmetric in ν and λ is presented in Sect. 6.

Eq. (4.8) exactly coincides with the large- N loop equation for the Hermitean one-matrix model with the logarithmic potential

$$\tilde{V}(\phi) = -(ab + c) \log(b - \phi) + (ab + c + 1) \log(a + \phi) - (a + b)\phi \quad (4.9)$$

for which

$$\tilde{V}'(\lambda) = \frac{a\lambda + c}{b - \lambda} + \frac{1 + c - b\lambda}{a + \lambda}. \quad (4.10)$$

The constant E_b is a free parameter which should be fixed requiring the analytic structure of E_λ on the complex λ -plane.

It is easy to demonstrate that the solution (4.6) satisfies Eq. (3.9). To show this let us use the symmetry of (4.6) in ν and λ and calculate $\text{Disc}_\nu G_{\lambda\nu}$ and $\text{Cont}_\nu G_{\lambda\nu}$ which read

$$\frac{1}{\pi\rho(\nu)} \text{Disc}_\nu G_{\lambda\nu} = \frac{\frac{a+\nu}{b-\lambda}}{\nu + E_\lambda - \frac{a\lambda+c}{b-\lambda}}, \quad \text{Cont}_\nu G_{\lambda\nu} = \frac{E_\lambda - \frac{\frac{1}{2}(a+\nu)\tilde{V}'(\nu)-1}{b-\lambda}}{\nu + E_\lambda - \frac{a\lambda+c}{b-\lambda}}. \quad (4.11)$$

These expressions identically satisfy Eq. (3.9) and, therefore, Eq. (3.10). I conclude that (4.6) solves the Riemann-Hilbert problem and should coincide with (3.11) for the potential $\mathcal{V}'$ given by Eq. (4.3). Analogously, Eq. (3.13) should reduce to Eq. (4.8).

It is instructive to discuss the one-cut solution to Eq. (4.8) which reads [17]

$$E_\lambda = \int_{C_1} \frac{d\omega}{4\pi i} \frac{\tilde{V}'(\omega)}{\lambda - \omega} \frac{\sqrt{(\lambda - x)(\lambda - y)}}{\sqrt{(\omega - x)(\omega - y)}} \quad (4.12)$$

where the ends of the cut, x and y , are determined by the asymptotic conditions

$$\int_{C_1} \frac{d\omega}{2\pi i} \frac{\tilde{V}'(\omega)}{\sqrt{(\omega - x)(\omega - y)}} = 0 \quad \int_{C_1} \frac{d\omega}{2\pi i} \frac{\omega \tilde{V}'(\omega)}{\sqrt{(\omega - x)(\omega - y)}} = 2. \quad (4.13)$$

For $\tilde{V}'$ given by (4.10) the contour integral can easily be calculated taking the residues at $\omega = \lambda, b$ and $-a$ while the residue at infinity vanishes since E_ω falls down as $1/\omega$. One gets

$$E_\lambda = \frac{1}{2} \left(\frac{a\lambda + c}{b - \lambda} + \frac{1 + c - b\lambda}{a + \lambda} \right) - \frac{1}{2} \left(\frac{ab + c}{b - \lambda} \frac{1}{\sqrt{(b - x)(b - y)}} + \frac{ab + c + 1}{a + \lambda} \frac{1}{\sqrt{(-a - x)(-a - y)}} \right) \sqrt{(\lambda - x)(\lambda - y)}. \quad (4.14)$$

It is worth noting that all the formulas recover the ones for the Gaussian potential in the limit (4.2). Since in the Gaussian case Eq. (4.13) has the solution

$$y = -x, \quad x^2 = \frac{a}{b} - \frac{b}{a}, \quad (\text{quadratic potential}), \quad (4.15)$$

the one-cut solution (4.14) is always realized for a and b which are big enough for the points b and $-a$ to lie outside of the cut $[x, y]$.

The situation when either b or $-a$ touch the end of the cut is associated with a phase transition to a more-than-one-cut solution. The one-matrix model with the potential (4.9) has at least as rich structure as that with the cubic potential since the latter can be obtained from (4.9) taking the limit

$$a, b \rightarrow \infty, \quad (a - b) \sim b^{\frac{1}{3}} \quad (\text{cubic potential}) \quad (4.16)$$

and rescaling the field $\phi \rightarrow \phi b^{\frac{1}{3}}$ which can always be done for the one-matrix model. The linear in ϕ term is finite providing $c \lesssim b^{\frac{2}{3}}$. It can be explicitly seen that in the limit (4.16) the one-cut (4.14) solution recovers the one for the cubic potential.

Since $\tilde{V}'(\lambda)$ is known, the function $F(\lambda)$ can be determined from Eq. (2.20) to be

$$F(\lambda) = \frac{b\lambda - c - 1}{a + \lambda}. \quad (4.17)$$

Now Eq. (2.6) determines

$$V'(\lambda) = \frac{a\lambda + c}{b - \lambda} + (2D - 1) \frac{b\lambda - c - 1}{a + \lambda} \quad (4.18)$$

which corresponds to the potential

$$V(\phi) = -(ab + c) \log(b - \phi) - (2D - 1)(ab + c + 1) \log(a + \phi) + [(2D - 1)b - a]\phi. \quad (4.19)$$

This formula recovers at $D = 1/2$ the potential (4.1) of the two-matrix model and at $D = 0$ the potential (4.9) of the associated one-matrix model. In the Gaussian limit (4.2) the solution found by Gross [11] is reproduced. Analogously, I verified that the next term of the $1/b$ expansion recovers a perturbation of the model with the quadratic potential by a cubic term with the coupling $g_3 = a/b^2$.

5 The “naive” continuum limit

The potential (4.19) of the D -dimensional Kazakov–Migdal model admits the following “naive” continuum limit. Let us expand V in ϕ which yields

$$\begin{aligned} V(\phi) = & \left(\frac{c}{b} - (2D - 1) \frac{c + 1}{a} \right) \phi + \frac{1}{2} \left(\frac{ab + c}{b^2} + (2D - 1) \frac{ab + c + 1}{a^2} \right) \phi^2 \\ & + \frac{1}{3} \left(\frac{ab + c}{b^3} - (2D - 1) \frac{ab + c + 1}{a^3} \right) \phi^3 \\ & + \frac{1}{4} \left(\frac{ab + c}{b^4} + (2D - 1) \frac{ab + c + 1}{a^4} \right) \phi^4 + \mathcal{O}(\phi^5). \end{aligned} \quad (5.1)$$

Now the idea is to choose the couplings a, b and c to provide the canonical scaling

$$\phi = \varepsilon^{\frac{D}{2}-1} \Phi, \quad V(\phi) = \varepsilon^D W(\Phi) \quad (5.2)$$

which is prescribed, as usual, by the kinetic term. The expansion will be justified choosing $a \sim b \rightarrow \infty$ so that $\phi/b \rightarrow 0$. The linear in ϕ term has the continuum limit providing

$$\left(\frac{a}{b} + 1 - 2D\right)c - (2D - 1) \lesssim \varepsilon^{\frac{D}{2}+1}a \quad (5.3)$$

which can always be satisfied by a proper choice of c .

The bare mass

$$m_0^2 = \frac{a}{b} + (2D - 1)\frac{b}{a} \geq 2D \quad (5.4)$$

for any D and the equality sign holds only for $D = 1$ at $a/b = 1$. Since the equality is needed to cancel the term $2D\phi^2$ coming from the kinetic energy, one concludes that the “naive” continuum limit is possible only for $D = 1$.

At $D = 1$ one has

$$m_0^2 = 2 + \frac{(a - b)^2}{ab} = 2 + \varepsilon^2 M^2 \quad (5.5)$$

where

$$\frac{a - b}{\varepsilon b} = M \sim 1 \quad (5.6)$$

is the continuum mass. Now the cubic in ϕ term has the continuum limit providing

$$a = b = \sqrt{\frac{2}{g}} \varepsilon^{\frac{D}{2}-2} \quad (5.7)$$

which justifies the expansion in

$$\frac{\phi}{b} \sim \varepsilon. \quad (5.8)$$

Notice that the quartic in ϕ term is also finite for the ε -dependence of b given by Eq. (5.7) as well as Eq. (5.3) is satisfied for any $c \sim 1$. We arrive, therefore, at the one-dimensional continuum quartic action

$$S[\Phi] = \int dt \left(\text{const.} \Phi(t) + \frac{1}{2} \dot{\Phi}^2(t) + \frac{M^2}{2} \Phi^2(t) + \sqrt{\frac{g}{2}} M \Phi^3(t) + \frac{g}{4} \Phi^4(t) \right) \quad (5.9)$$

where $\dot{\Phi} \equiv \partial_t \Phi$. The coefficient in front of the linear in ϕ term can be made arbitrary (or vanishing) by choosing c .

The above procedure of taking the “naive” continuum limit works for $D < 4$ where b given by Eq. (5.7) is divergent. The only difference is that for $D \neq 1$ the bare mass term remains infinite. One can cancel it, however, for $2 \leq D < 4$ by the standard *renormalization* procedure. The restriction $D < 4$ is precisely the one where the scalar theory with the quartic interaction is renormalizable.

As was shown by Gross [11], any solution of the master field equation (3.13) for an arbitrary potential reproduces in $D = 1$ the spectral density given by the fermionic solution of Ref. [18]. For this reason there is no doubt that the solution of the previous section does. However, an interesting question is how the one-cut solution (4.14) can reproduce this $D = 1$ spectral density? The answer is that the potential (5.9) is simple enough to provide at $D = 1$ the factorized structure

$$\pi\rho(\lambda) = \sqrt{2E_F - 2W(\lambda)} = \sqrt{\frac{g}{2}}(\lambda + \text{const.})\sqrt{(y - \lambda)(\lambda - x)} \quad (5.10)$$

by turning the coupling in front of the linear in ϕ term.

6 The Itzykson–Zuber correlator

To calculate the pair correlator of U and $U^\dagger$, one should take the discontinuity of (4.6) both in ν and λ across the cut (cuts) [9]. For the solution (4.6) one gets

$$C(\nu, \lambda) \equiv \frac{1}{\pi^2 \rho(\nu) \rho(\lambda)} \text{Disc}_\lambda \text{Disc}_\nu G_{\nu\lambda} = \frac{(a + \nu)(a + \lambda)}{\mathcal{D}(\nu, \lambda)}, \quad \nu, \lambda \in \text{cut} \quad (6.1)$$

with

$$\begin{aligned} \mathcal{D}(\nu, \lambda) = & -\lambda^2 \nu^2 + (b - a)\lambda\nu(\lambda + \nu) + ab(\lambda^2 + \nu^2) - \lambda\nu(a^2 + b^2 + 2c + 1) \\ & + (\lambda + \nu)(b - ac + bc) - c^2 - b^2 + (ab + c)[(a + b)E_b - 1], \end{aligned} \quad (6.2)$$

where E_b depends on the type of the solution of Eq. (4.8) (one-cut or more-than-one-cut solutions). In the Gaussian limit (4.2) when $E_b \rightarrow 1/b$, one recovers the result of Ref. [9].

While the expression (6.1) looks complicated, the denominator seems to be positive. To see this let us rewrite $\mathcal{D}(\nu, \lambda)$ as

$$\mathcal{D}(\nu, \lambda) = (b - \nu)(a + \nu)(\lambda - r_+(\nu))(\lambda - r_-(\nu)) \quad (6.3)$$

and

$$r_\pm(\nu) = \frac{1}{2}(F(\nu) + \mathcal{V}'(\nu)) \pm i\pi\rho(\nu). \quad (6.4)$$

Since $\rho(\nu)$ is real for $\nu \in \text{cut}$, the roots are complex except for the values of parameters when $\rho(\nu)$ vanishes for some $\nu \in \text{cut}$. This might happen for the cases of phase transitions in the one-matrix model with the potential (4.9). The simplest one is from the one-cut solution to a two-cut solution. It is an algebraic problem to study whether the roots (6.4) always lie outside of the cut (cuts).

These phase transitions does not change, however, the functional structure of $C(\nu, \lambda)$ given by Eqs. (6.1), (6.2) — only the constant E_b changes. For this reason the Kazakov–Migdal model with the logarithmic potential (4.19) seems to be too simple to exhibit a non-trivial critical behavior at $D > 1$.

Acknowledgements

I am grateful to J. Ambjørn, D. Boulatov, M. Dobroliubov, C. Kristjansen and G. Semenoff for useful discussions. I thank the NBI high energy theory group for the hospitality at Copenhagen.

References

- [1] R.C. Penner, *Comm. Math. Phys.* **113** (1987) 299; *J. Diff. Geom.* **27** (1988) 35.
- [2] J. Distler and C. Vafa, *Mod. Phys. Lett.* **A6** (1991) 259.
- [3] C.-I Tan, *Mod. Phys. Lett.* **A6** (1991) 1373; *Phys. Rev.* **D45** (1992) 2862;
S. Chaudhuri, H. Dykstra and J. Lykken, *Mod. Phys. Lett.* **A6** (1991) 1665;
G. Gilbert and M. Perry, *Nucl. Phys.* **B364** (1991) 734.
- [4] L. Chekhov and Yu. Makeenko, *Phys. Lett.* **278B** (1992) 271;
S. Kharchev, A. Marshakov, A. Mironov and A. Morozov, *Nucl. Phys.* **B397** (1993) 339;
R. Dijkgraaf, G. Moore and R. Plesser, *Nucl. Phys.* **B394** (1993) 356.
- [5] J.-M. Daul, V.A. Kazakov and I.K. Kostov, *Rational theories of 2d gravity from the two-matrix model*, preprint CERN-TH-6834/93 (March, 1993).
- [6] M. Douglas, in *Random surfaces and quantum gravity (Cargèse 1990)*, eds. O. Alvarez *et al.*, Plenum, 1991, p.77.
- [7] V.A. Kazakov and A.A. Migdal, *Nucl. Phys.* **B397** (1993) 214.
- [8] V.A. Kazakov, *Nucl. Phys. B (Proc. Suppl.)* **30** (1993) 149.
- [9] M.I. Dobroliubov, Yu. Makeenko and G.W. Semenoff, *Correlators of the Kazakov-Migdal model*, UBC preprint (April, 1993).
- [10] A.A. Migdal, *Mod. Phys. Lett.* **A8** (1993) 359.
- [11] D. Gross, *Phys. Lett.* **293B** (1992) 181.
- [12] A.A. Migdal, *Mod. Phys. Lett.* **A8** (1993) 245.
- [13] Yu. Makeenko, *Matrix models of induced large- N QCD*, preprint ITEP-YM-11-92 (December, 1992).
- [14] M. Staudacher, *Phys. Lett.* **305B** (1993) 332.
- [15] J. Ambjørn and C. Kristjansen, unpublished.
- [16] D. Boulatov, unpublished.
- [17] A.A. Migdal, *Phys. Rep.* **102** (1983) 199.
- [18] E. Brézin, C. Itzykson, G. Parisi and J.B. Zuber, *Commun. Math. Phys.* **59** (1978) 35.